



# Enroflox<sup>™</sup> 100

(enrofloxacin)

## BEEF PRODUCERS ASKED FOR IT, NORBROOK DELIVERS...

### Introducing New Enroflox<sup>™</sup> 100 The Newest Addition To Your Arsenal For Treating BRD

- In cattle, for multi-day use only
- Same active ingredient as Baytril<sup>®</sup> 100 Injection
- Same formulation as Baytril 100 Injection
- FDA-Approved for the treatment of Bovine Respiratory Disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*

### Enroflox 100 Injection ... The **NEW** Choice

For use by or on the order of a licensed veterinarian. Enroflox100 is not approved for a one-day, single dose of therapy in cattle. Federal law prohibits the off-label use of this drug in food producing animals. Cattle intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal. Use with caution in animals with known or suspected CNS disorders. Observe label directions and withdrawal times. See product labeling for full product information.



Enroflox<sup>™</sup> 100  
(enrofloxacin)



FOR VETERINARY USE ONLY

[www.norbrookinc.com](http://www.norbrookinc.com)

The Norbrook logos are registered trademarks of Norbrook Laboratories Limited  
Enroflox is a trademark of Norbrook Laboratories Limited  
Baytril is a registered trademark of Bayer Animal Health



**Norbrook<sup>®</sup>**

## DOSAGE ADMINISTRATION

- **Enroflox 100** is administered as an SQ, **multiple-day therapy**
  - Selection of the appropriate dose and duration of therapy should be based on an assessment of the severity of disease, pathogen susceptibility and clinical response
  - Administer daily, an SQ dose of 2.5 - 5.0 mg/kg of body weight (1.1 - 2.3 mL/100 lb.)
  - Treatment should be repeated at 24-hour intervals for three days
  - Additional treatments may be given on days 4 and 5 to animals that have shown clinical improvement but not total recovery
- **Enroflox 100** dose volume should not exceed 20 mL per injection site
- **Enroflox 100** contains 100 mg of enrofloxacin per mL



Cattle Weight (lbs.)	Multiple-Day Therapy 2.5 - 5.0 mg/kg Dose Size (mL)
100	1.5 - 2.0
200	2.5 - 4.5
300	3.5 - 6.5
400	4.5 - 9.0
500	5.5 - 11.5
600	7.0 - 13.5
700	8.0 - 16.0
800	9.0 - 18.0
900	10.0 - 20.5
1000	11.0 - 23.0
1100	12.5 - 25.0

## PRODUCT DESCRIPTION

- **Enroflox 100** is a FDA-approved sterile, ready-to-use injectable antimicrobial solution that contains enrofloxacin, a broad-spectrum fluoroquinolone antimicrobial agent
- **Enroflox 100** is administered as a multiple-day therapy
- **Enroflox 100** is labeled for the treatment of Bovine Respiratory Disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*, major pathogenic bacteria associated with BRD
- *In vitro*<sup>\*</sup>, enrofloxacin kills 97% of BRD-causing bacteria in 1-2 hours<sup>1,2</sup>



## PRODUCT AVAILABILITY

**Enroflox 100** is available in convenient 100 mL and 250 mL doses to fit any size operation.

Contact your veterinarian today for more information or to purchase new **Enroflox 100**.

**Enroflox™ 100 Injection ... The *NEW* Choice**

**Enroflox™ 100**  
(enrofloxacin)

<sup>\*</sup> The clinical significance of *in vitro* data has not been demonstrated.

<sup>1</sup> Blondeau J.M., Borsos S., Blondeau L.D., Blondeau B.J., Hesje C. The killing of clinical isolates of *Mannheimia haemolytica* (MH) by enrofloxacin (ENR) using minimum inhibitory and mutant prevention drug concentrations and over a range of bacterial inocula. In: *ASM Conference on Pasteurellaceae*; 2005 October 23-26; Kohala Coast, Big Island, Hawaii: American Society of Microbiology; 2005. Abstract B12.

<sup>2</sup> Blondeau J.M., Borsos S.D., Hesje C.H., Blondeau L.D., Blondeau B.J. Comparative Killing of Bovine Isolates of *Mannheimia haemolytica* by Enrofloxacin, Florfenicol, Tilmicosin and Tulathromycin Using the Measured Minimum Inhibitory Concentration (MIC) and Mutant Prevention Concentration (MPC) Drug Values. In: *International Meeting of Emerging Diseases and Surveillance (IMED)*, Vienna, Austria, February 23-25, 2007. Figures 8-10.

# Enrofloxacin (enrofloxacin)

100 mg/mL Antimicrobial  
Injectable Solution



For Subcutaneous Use in Beef Cattle, Non-Lactating Dairy Cattle and Swine Only.  
Not for Use in Female Dairy Cattle 20 Months of Age or Older Or in Calves To Be Processed For Veal.


**CAUTION:**  
Federal (U.S.A.) law restricts this drug to use by or on the order of a licensed veterinarian.  
Federal (U.S.A.) law prohibits the extra-label use of this drug in food producing animals.

**PRODUCT DESCRIPTION:**

Enroflox 100 is a sterile, ready-to-use injectable antimicrobial solution that contains enrofloxacin, a broad-spectrum fluoroquinolone antimicrobial agent. Therapeutic treatment with Enroflox 100 is administered as a single dose for one day (swine) or for multiple days (cattle) of therapy. Each mL of Enroflox 100 contains 100 mg of enrofloxacin. Excipients are L-arginine base 200 mg, n-butyl alcohol 30 mg, benzyl alcohol (as a preservative) 20 mg and water for injection q.s.

**CHEMICAL NOMENCLATURE AND STRUCTURE:**

1-cyclopentyl-7-(4-ethyl-1-piperazinyl)-6-fluoro-1,4-dihydro-4-oxo-3-quinolonecarboxylic acid.



**INDICATIONS:**

**Cattle:** Enroflox 100 is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* in beef and non-lactating dairy cattle.

**Swine:** Enroflox 100 is indicated for the treatment and control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Haemophilus parasuis* and *Streptococcus suis*.

**DOSEAGE AND ADMINISTRATION:**

Enroflox 100 is administered as a single dose for one day (swine) or for multiple days (cattle) of therapy.

**Cattle:** Administer daily a subcutaneous dose of 2.5 – 5.0 mg/kg of body weight (1.1 – 2.3 mL/100 lb). Treatment should be repeated at 24-hour intervals for three days. Additional treatments may be given on Days 4 and 5 to animals that have shown clinical improvement but not total recovery. Administered dose volume should not exceed 20 mL per injection site.

Table 1 – Enroflox 100 Dose and Treatment Schedule for Cattle\*

WEIGHT (lb)	Multiple-Day Therapy** 2.5 - 5.0 mg/kg Dose Volume (mL)
100	1.5 - 2.0
200	2.5 - 4.5
300	3.5 - 6.5
400	4.5 - 9.0
500	5.5 - 11.5
600	7.0 - 13.5
700	8.0 - 16.0
800	9.0 - 18.0
900	10.0 - 20.5
1000	11.0 - 23.0
1100	12.5 - 25.0

\*Dose volumes have been rounded to the nearest 0.5 mL within the dose range.

\*\*Not for Single-Dose Therapy.

**Swine:** Administer once behind the ear, a subcutaneous dose of 7.5 mg/kg of body weight (3.4 mL/100 lb). Administered dose volume should not exceed 5 mL per injection site.

Table 2 – Enroflox 100 Dose and Treatment Schedule for Swine

WEIGHT (lb)	Dose Volume (mL)
50	1.7
100	3.4
150	5.1
200	6.8
250	8.5

**RESIDUE WARNINGS:**

**Cattle:** Animals intended for human consumption must not be slaughtered within 28 days from the last treatment. This product is not approved for female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Swine:** Animals intended for human consumption must not be slaughtered within 5 days of receiving a single-injection dose.

**HUMAN WARNINGS:**

**Keep out of the reach of children.**  
Avoid contact with eyes. In case of contact, immediately flush eyes with copious amounts of water for 15 minutes. In case of dermal contact, wash skin with soap and water. Consult a physician if irritation persists following ocular or dermal exposures. Individuals with a history of hypersensitivity to quinolones should avoid this product. In humans, there is a risk of user photosensitization within a few hours after excessive exposure to quinolones. If excessive accidental exposure occurs, avoid direct sunlight. For customer service, to obtain a copy of the Material Safety Data Sheet (MSDS) or to report adverse reactions, call Norbrook at 1-866-591-5777.

**PRECAUTIONS:**

The effects of enrofloxacin on cattle or swine reproductive performance, pregnancy and lactation have not been adequately determined.  
The long-term effects on articular joint cartilage have not been determined in pigs above market weight.

Subcutaneous injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter. Enroflox 100 contains different excipients than other enrofloxacin products. The safety and efficacy of this formulation in species other than cattle and swine have not been determined.

Quinolone-class drugs should be used with caution in animals with known or suspected Central Nervous System (CNS) disorders. In such animals, quinolones have, in rare instances, been associated with CNS stimulation which may lead to convulsive seizures. Quinolone-class drugs have been shown to produce erosions of cartilage of weight-bearing joints and other signs of arthropathy in immature animals of various species. See Animal Safety section for additional information.

**ADVERSE REACTIONS:**

No adverse reactions were observed during clinical trials.

**MICROBIOLOGY:**

Enrofloxacin is bactericidal and exerts its antibacterial effect by inhibiting bacterial DNA gyrase (a type II topoisomerase) thereby preventing DNA supercoiling and replication which leads to cell death. Enrofloxacin is active against Gram-negative and Gram-positive bacteria.

**EFFECTIVENESS:**

**Cattle:** A total of 845 calves with naturally-occurring BRD were treated with enrofloxacin injection in eight field trials located in five cattle-feeding states. Response to treatment was compared to non-treated controls. BRD and mortality were significantly reduced in enrofloxacin-treated calves. No adverse reactions were reported in treated animals.

**Swine:** A total of 590 pigs were treated with enrofloxacin injection or saline in two separate natural infection SRD field trials. For the treatment of SRD, the success rate of enrofloxacin-treated pigs that were defined as sick and febrile (increased respiratory rate, labored or dyspneic breathing, depressed attitude and a rectal temperature  $\geq 104.0^{\circ}\text{F}$ ) was statistically significantly greater than the success rate of saline-treated "sick and febrile" pigs. For the control of SRD, mean rectal temperature, mortality (one trial) and morbidity were statistically significantly lower for enrofloxacin-treated pigs in pens containing a percentage of "sick and febrile" pigs compared to saline-treated pigs.

**TOXICOLOGY:**

The oral LD50 for laboratory rats was greater than 5000 mg/kg of body weight. Niney-day feeding studies in dogs and rats revealed no observable adverse effects at treatment rates of 3 and 40 mg/kg respectively. Chronic studies in rats and mice revealed no observable adverse effects at 5.3 and 52 mg/kg respectively. There was no evidence of carcinogenic effect in laboratory animal models. A two-generation rat reproduction study revealed no effect with 10 mg/kg treatment. No teratogenic effects were observed in rabbits at doses of 25 mg/kg or in rats at 50 mg/kg.

**ANIMAL SAFETY:**

**Cattle:** Safety studies were conducted in feeder calves using single doses of 5, 15, and 25 mg/kg for 15 consecutive days and 50 mg/kg for 5 consecutive days. No clinical signs of toxicity were observed when a dose of 5 mg/kg was administered for 15 days. Clinical signs of depression, incoordination, and muscle fasciculation were observed in calves when doses of

15 or 25 mg/kg were administered for 10 to 15 days. Clinical signs of depression, inappetence and incoordination were observed when a dose of 50 mg/kg was administered for 3 days. No drug-related abnormalities in clinical pathology parameters were identified. No articular cartilage lesions were observed after examination of stiffler joints from animals administered 25 mg/kg for 15 days.

A safety study was conducted in 23-day-old calves using doses of 5, 15, and 25 mg/kg for 15 consecutive days. No clinical signs of toxicity or changes in clinical pathology parameters were observed. No articular cartilage lesions were observed in the stiffler joints at any dose level at 7 days and 9 days following 15 days of drug administration.

An injection site study conducted in feeder calves demonstrated that the formulation may induce transient reaction in the subcutaneous tissue and underlying muscle. No painful responses to administration were observed.

**Swine:** A safety study was conducted in 32 pigs weighing approximately 57 kg (125 lb) using single doses of 5, 15, or 25 mg/kg daily for 15 consecutive days. Incidental lameness of short duration was observed in all groups, including the saline-treated controls. Musculoskeletal stiffness was observed following the 15 and 25 mg/kg treatments with clinical signs appearing during the second week of treatment. Clinical signs of lameness improved after treatment ceased and most animals were clinically normal at necropsy.

A second study was conducted in two pigs weighing approximately 23 kg (50 lb), treated with 50 mg/kg for 5 consecutive days. There were no clinical signs of toxicity or pathological changes.

An injection site study conducted in pigs demonstrated that the formulation may induce a transient reaction in the subcutaneous tissue. No painful responses to administration were observed.

**STORAGE CONDITIONS:**

Protect from direct sunlight. Do not refrigerate or freeze. Store below 77°F (25°C). Precipitation may occur due to cold temperature. To redissolve, warm and then shake the vial.

**HOW SUPPLIED:**

Enroflox 100: 100 mL Bottle  
100 mg/mL  
250 mL Bottle  
100 mg/mL

**REFERENCES:**

1. Hooper, D. C., Wolfson, J. S., *Quinolone Antimicrobial Agents*, 2nd ed., 59 - 75, 1993.

For customer service, to obtain a copy of the Material Safety Data Sheet (MSDS) or to report adverse reactions, call Norbrook at 1-866-591-5777.

**Restricted Drug - California. Use Only as Directed.**

Made in the UK

Norbrook Laboratories Limited  
Newry, BT35 6PU, Co. Down,  
Northern Ireland

05167007



Please  
Contact  
Us At:



53 Industrial Circle  
Lancaster, PA 17601

Phone: 1-800-233-0210  
Online: www.pennvet.com